

Bulk Gauge Fields in AdS Supergravity and Supersingletons

Sergio Ferrara

CERN Geneva, Switzerland

`Sergio.Ferrara@cern.ch`

Alberto Zaffaroni

CERN Geneva, Switzerland

`Alberto.Zaffaroni@cern.ch`

We describe conformal operators living at the boundary of AdS_{d+1} in a general setting. Primary conformal operators at the threshold of the unitarity bounds of UIR's of $O(d,2)$ correspond to singletons and massless fields in AdS_{d+1} , respectively. For maximal supersymmetric theories in AdS_{d+1} we describe “chiral” primary short supermultiplets and non-chiral primary long supermultiplets. Examples are exhibited which correspond to KK and string states. We give the general contribution of a primary conformal operator to the OPE and Green's functions of primary fields, which may be relevant to compute string corrections to the four-point supergraviton amplitude in Anti-de-Sitter space.

The material in this paper was presented by one of the authors (S.F) at String 98, Santa Barbara.

CERN-TH/98-229

July 98

Some aspects of the $\text{AdS}_{d+1}/\text{CFT}$ correspondence, inspired by the original conjecture by Maldacena[1], and subsequently sharpened by Gubser, Klebanov, Polyakov and Witten [2,3], are considered in this paper¹. The maximal supersymmetric case, corresponding to the horizon geometry of D3 branes in type IIB strings, is considered as basic example.

More specifically, we describe the relation between interacting non-abelian singletons on ∂AdS_5 , i.e. $\text{N}=4$ $\text{SU}(n)$ super Yang-Mills theory and the bulk supergravity theory on $\text{AdS}_5 \times S^5$.

In section 1, we review some properties of primary conformal fields on ∂AdS_5 and, in section 2, their superextension. In sections 3 and 4, we describe several types of $\text{N}=4$ multiplets, realized in $\text{N}=4$ super Yang-Mills theory, which in the AdS/CFT correspondence, are related to singletons, massless and massive KK chiral primary supermultiplets as well as massive higher spin multiplets. In section 5, we briefly describe manifestly covariant OPE on ∂AdS_5 which can be used to compute boundary correlation functions of primary operators corresponding to states in the spectrum of the type IIB string compactified on $\text{AdS}_5 \times S^5$. In particular, the contribution to the four-point function of chiral primaries, due to the non-chiral primaries corresponding to string states, is exhibited. The latter may be used to connect string corrections to the four-point graviton amplitude in $\text{AdS}_5 \times S^5$ geometry to four-point stress-energy tensor correlation functions on the boundary.

1. Conformal fields on ∂AdS_5

The 15 generators $J_{AB} = -J_{BA}$ of the conformal algebra $\text{O}(4,2)$ can be defined in terms of the Poincaré generators $P_\mu, M_{\mu\nu}$, the dilatation D and the special conformal transformation K_μ , by the relations,

$$J_{\mu\nu} = M_{\mu\nu}, \quad J_{5\mu} = \frac{1}{2}(P_\mu - K_\mu), \quad J_{6\mu} = \frac{1}{2}(P_\mu + K_\mu), \quad J_{65} = D. \quad (1.1)$$

The irreducible representations of the conformal algebra are specified by the values of the three Casimir operators [5],

$$\begin{aligned} C_I &= J^{AB} J_{AB} = \\ C_{II} &= \epsilon_{ABCDE F} J^{AB} J^{CD} J^{EF} \\ C_{III} &= J_A^B J_B^C J_C^D J_D^A \end{aligned} \quad (1.2)$$

¹ For earlier work on the connection between branes geometry, Anti-de-Sitter space and singletons, see [4]

Every irreducible (infinite dimensional) representation can be specified by an irreducible representation of the Lorentz group with definite conformal dimension and annihilated by K_μ . In terms of the stability algebra at $x = 0$ ($K_\mu, D, M_{\mu\nu}$), we define a primary conformal field, in a generic representation of the Lorentz group, by

$$\begin{aligned} [O_{\{\alpha\}}(0), D] &= i l O_{\{\alpha\}}(0) \\ [O_{\{\alpha\}}(0), K_\mu] &= 0. \end{aligned} \tag{1.3}$$

The descendents $\partial \dots \partial O_{\{\alpha\}}(0)$ fill an infinite dimensional representation specified by three numbers (l, j_L, j_R) , which contain the conformal dimension and the Lorentz quantum number of the primary operator.

In terms of the three Casimirs of the stability algebra,

$$D = l, \quad \frac{1}{2} M_{\mu\nu} M^{\mu\nu} = j_L(j_L + 1) + j_R(j_R + 1), \quad \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} M^{\mu\nu} M^{\rho\sigma} = j_L(j_L + 1) - j_R(j_R + 1) \tag{1.4}$$

the $O(4,2)$ Casimirs take the following values[6]:

$$\begin{aligned} C_I &= l(l - 4) + 2j_L(j_L + 1) + 2j_R(j_R + 1) \\ C_{II} &= (l - 2)(j_L(j_L + 1) - j_R(j_R + 1)) \\ C_{III} &= (l - 2)^4 - 4(l - 2)^2[j_L(j_L + 1) + j_R(j_R + 1) + 1] + 16j_L j_R(j_L + 1)(j_R + 1) \end{aligned} \tag{1.5}$$

In the particular case of a tensor representation of spin s ($l, j_L = j_R = s/2$), we have:

$$\begin{aligned} C_I &= l(l - 4) + s(s + 2) \\ C_{II} &= 0 \\ C_{III} &= [l(l - 2) - s(s + 2)][(l - 2)(l - 4) - s(s + 2)] \end{aligned} \tag{1.6}$$

Let us consider, for example, the case of a conformal scalar. It is convenient to consider a six-dimensional space with signature $(+ - - -, - +)$ where the $O(4,2)$ generators act on the coordinates $\eta_A, A = 0, \dots, 5$ as $L_{AB} = i(\eta_A \partial_B - \eta_B \partial_A)$ [7,5]. The conformal compactification of Minkowski space-time can be identified with the hypercone $\eta_A \eta^A = 0$, with projectively identified coordinates ($\eta^A = \lambda \eta^A$). A conformal scalar can be represented as a homogeneous function on the hypercone $\eta_A \eta^A = 0$ [7,5],

$$(\eta \partial) \Phi = \lambda \Phi, \quad (\lambda = -l) \tag{1.7}$$

The quadratic Casimir

$$\frac{1}{2}L_{AB}L^{AB} = -\eta^2\partial^2 + \eta\partial(4 + \eta\partial) \quad (1.8)$$

on the hypercone reduces to $C_I = l(l - 4)$.

To describe tensor conformal fields, we can consider homogeneous tensors $O_{A_1\dots A_s}(\eta)$ on the hypercone. The irreducible representations of the conformal algebra are specified by the symmetric-traceless tensors which also satisfy refs[5],[8],

$$\begin{aligned} \eta^{A_1}O_{A_1\dots A_s}(\eta) &= 0 \\ \partial^{A_1}O_{A_1\dots A_s}(\eta) &= 0 \end{aligned} \quad (1.9)$$

Acting on these tensors, the Casimirs are not purely orbital, but they must be supplemented with an internal part, corresponding to a finite dimensional representation of $O(4,2)$, $L_{AB}\delta_{A_1\dots A_s}^{B_1\dots B_s} + \Sigma_{A_1\dots A_s}^{B_1\dots B_s}$.

Unitarity imposes the following bounds [9],

$$l \geq 1 + j \quad (j_L j_R = 0) \quad (1.10)$$

$$l \geq 2 + j_L + j_R \quad (j_L j_R \neq 0) \quad (1.11)$$

The two unitarity thresholds are satisfied by massless fields and conserved tensors field, respectively [8]. The equations

$$\partial^2\Phi_{(0,j)} = 0 \quad (1.12)$$

$$\partial^{\alpha_1\dot{\alpha}_1}O_{\alpha_1\dots\alpha_{2j_L},\dot{\alpha}_1\dots\dot{\alpha}_{2j_R}} = 0 \quad (1.13)$$

are indeed conformal covariant only if $l = 1 + j$ and $l = 2 + j_L + j_R$, respectively. This can be easily proved by considering the $O(4,2)$ commutation rule $[K_\mu, P_\mu] = -2i(g_{\mu\nu}D + M_{\mu\nu})$.

In the AdS_5/CFT correspondence, all gauge invariant composite operators in the CFT can be associated with fields in AdS_5 [2,9,3]. $O(4,2)$ is reinterpreted as the isometry group of AdS_5 , and particles in AdS_5 are classified by the quantum number (E_0, j_L, j_R) of the maximal compact subgroup $O(2) \times O(4)$. In the identification with CFT operators, E_0 is identified with the scaling dimension and (j_L, j_R) with the 4-dimensional Lorentz quantum number of the primary conformal operator.

The $O(4,2)$ covariant wave equation for a particle with quantum numbers (E_0, j_L, j_R) in AdS_5 can be expressed in terms of the Casimirs of the conformal algebra. In this way, the mass of a particle can be expressed as a function of (E_0, j_L, j_R) . In the case of a scalar,

for example, the Laplace operator in AdS_5 coincides with the quadratic Casimir C_I and the wave equation reads,

$$\partial^2 \Phi - \eta \partial(4 + \eta \partial) = -\frac{1}{2} L_{AB} L^{AB} \Phi = E_0(E_0 - 4) \Phi, \quad (1.14)$$

where we used eq. (1.6) with $l \rightarrow E_0$. We see that the mass square of a scalar field in AdS_5 can be expressed in terms of the conformal dimension E_0 of the field by $m^2 = E_0(E_0 - 4)$.

In general we obtain the following relations [10]:

– Scalars: ϕ

$$D(E_0, 0, 0) : \quad m^2 = C_I = E_0(E_0 - 4) \quad (1.15)$$

– Vector field: A_μ

$$D\left(E_0, \frac{1}{2}, \frac{1}{2}\right) : \quad m^2 = C_I = (E_0 - 1)(E_0 - 3) \quad (1.16)$$

– Symmetric tensor: $g_{\mu\nu} = g_{\nu\mu}$

$$D(E_0, 1, 1) : \quad m^2 = C_I - 8 = E_0(E_0 - 4) \quad (1.17)$$

– Antisymmetric tensor: $A_{\mu\nu} = -A_{\nu\mu}$

$$D(E_0, 1, 0) \oplus D(E_0, 0, 1) : \quad m^2 = C_I = (E_0 - 2)^2 \quad (1.18)$$

and for fermions,

– Fermions of spin 3/2: ψ_μ

$$D\left(E_0, 1, \frac{1}{2}\right) + D\left(E_0, \frac{1}{2}, 1\right) ; \quad m = E_0 - 2 \quad (1.19)$$

– Fermions of spin 1/2: λ

$$D\left(E_0, 0, \frac{1}{2}\right) + D\left(E_0, \frac{1}{2}, 0\right) ; \quad m = E_0 - 2 \quad (1.20)$$

Of particular relevance also in AdS_5 are the representations of $\text{O}(4,2)$ which saturate the unitarity bounds. The states which saturate the bound (1.10) in AdS_5 are called singletons and are topological fields living at the boundary of AdS_5 [11,12,13]. They cannot be associated with any gauge invariant operator in the CFT, but it is suggestive that they have the same quantum number of the conformal (generally colored) fundamental fields appearing in the CFT. In particular, in the well known case of the duality between type

IIB on $\text{AdS}_5 \times S^5$ and N=4 SYM, the singletons in AdS_5 are in correspondence with the fundamental N=4 SYM multiplet. All the other unitary representations in AdS_5 propagate inside the bulk of AdS_5 and are in correspondence with gauge invariant composite operators in the CFT.

Let us discuss also the meaning of the second unitarity bound in AdS (1.11). This is the case of conserved currents in the CFT. The associated fields in AdS_5 are massless fields sustaining a gauge invariance. In this way, the global symmetries in CFT are associated with local symmetries in AdS_5 [9]. Here are the simplest examples of this correspondence:

- stress-energy tensor/graviton $T_{\mu\nu} \rightarrow g_{\mu\nu}$
- global current/gauge field $J_\mu \rightarrow A_\mu$
- supercurrent/gravitino $J_{\mu\alpha} \rightarrow \Psi_{\mu\alpha}$.

The conserved tensors in this list satisfy in CFT an equation like (1.13). This guarantees that the number of degrees of freedom contained in a conserved tensor matches with those of a massless particle in AdS_5 . The number of degrees of freedom of a conserved tensor is $(2j_L + 1)(2j_R + 1) - 4j_L j_R = 2(j_L + j_R) + 1$, which can exactly sustain a representation of spin $(j_L + j_R)$ of the little group $O(3)$ of massless particles in the Poincaré limit of AdS_5 .

In the case of representations for which $l > 2 + j_L + j_R$, the fields in AdS_5 are massive, and the corresponding CFT operators are not conserved.

2. The maximal supersymmetric case

In the AdS/CFT correspondence, the conformal dimension E_0 is not in general an integer. Only the conserved currents, in general, are protected under renormalization and have integer conformal dimensions; they are indeed associated with massless states in AdS. On the other hand, the generic CFT operator is expected to have anomalous dimensions. Introducing supersymmetry in the game, we have the notion of a “chiral” primary conformal operator, whose dimension is not renormalized even if it is not a conserved tensor. In the familiar case of N=1 theories, the non-renormalization of the conformal dimension follows from the relation with the $U(1)_R$ charge ($E_0 = q$). The notion of chirality is associated with a shortening of the supersymmetric multiplet.

The Maldacena’s conjecture [1] relates a SCFT in d spacetime dimensions with N extended supersymmetries with the type IIB string (or M theory) on $\text{AdS}_{d+1} \times H$, where H is an Einstein manifold which gives rise, after dimensional reduction, to a $2N$ extended gauged supergravity in AdS_{d+1} . Let us focus on the maximal supersymmetric case for $d=3$

or $d=4$. The conformal groups $O(3,2)$ and $O(4,2)$ are enlarged to the supergroups $O(8/4)$ and $SU(2,2/4)$, with 32 conformal supercharges.

The superconformal algebra representations which saturate the bound (1.10) correspond to the AdS_{d+1} singletons, which are associated with the fundamental massless conformal fields defining the SCFT. The superconformal algebra representations which saturate the bound (1.11) correspond to the *massless* multiplets in AdS_{d+1} , associated with the CFT multiplet of global currents. Other fields besides the conserved tensors are, in general, required to close a supersymmetric multiplet²; in the same way, the AdS_{d+1} *massless* multiplet contains some massive fields, in addition to the massless graviton, gravitinos and gauge fields [16]. These massless multiplets are automatically short and their dimension is protected. The massive multiplets can be short (chiral, with canonical dimensions) or long (with anomalous dimensions). From the supergravity side, we know that all the KK states, coming from the dimensional reduction on H , are in short representations [17] and have integer dimensions. On the other hand, a generic string state has a mass which is not an integer and the corresponding CFT operator acquires anomalous dimension (in generally very large, in the limit in which supergravity can be trusted) [2,3].

The superconformal representations can be induced by conformal primary fields. A generic scalar superfield has 2^{16} components with spin range from 0 up to 4. The degeneracy of representations of the Lorentz group $O(d-1,1)$ is $Sp(16)$ for $d=3$ and $Sp(8)$ for $d=4$. Generic superfields correspond to representations of the Clifford algebra of $O(32)$ where left and right representations are the bosons and fermions, respectively.

Chiral primary superfields have $2^8 \times l$ components, where l is the (finite) dimension of some representation of the Lorentz group and the R-symmetry ($O(8)$ for $d=3$, $SU(4)$ for $d=4$).

Let us now focus on $d=4$. An unconstrained superfields, with θ_α^A in the $(1/2,0)$ representation of $SL(2,C)$ and N of $SU(N)$, has generically 2^{4N} components, spanning the Clifford algebra of $SO(4N)$ [15]. Bosons and fermions, corresponding to the even and odd powers in the θ 's expansion, are the two chiral spinorial representations of $O(4N)$. These superfields are extended to a representation of $SU(2,2/N)$.

² For example, three set of scalars in different representation of $SU(4)$, fermions and tensor fields are required to close the $N=4$ SYM supercurrent multiplet [14,15] which contains the conserved stress-energy tensor, the spinorial supersymmetry currents and the $SU(4)$ R-symmetry currents.

There are three types of different supermultiplets.

The ultrashort representations correspond to singleton representations. These multiplets have degeneracy $2^4(2j_L + 1)$ with spin range from $j_L - 1$ to $j_L + 1$ [6]. They correspond to a sequence of massless conformal fields with dimension and spin related as in eq. (1.10). The N=4 SYM fundamental multiplet belongs to this class of representations, and it is special because it is self-conjugated.

Short representations correspond to massless or massive representations with degeneracy $2^8 r$ where r is some finite dimensional representation of $SL(2, C) \times SU(4)$. Generic massless representations with spin range from $(j_L - 1, j_R - 1)$ to $(j_L + 1, j_R + 1)$ can be obtained by tensoring two singleton representations $(0, j_L) \times (j_R, 0)$ giving $2^8(2j_L + 1)(2j_R + 1)$ states [6]. The physical sector of these massless representations is obtained by having a gauge symmetry in AdS_5 , which reduces the number of components to $2^8(2(j_L + j_R) + 1)$ [18,9,10]. These massless representations are obtained by a sequence of transverse conformal primary fields with dimension and spin satisfying eq. (1.11). Generically, a N-extended superfield corresponding to a short multiplet has an expansion in half of the θ 's. It can be chiral or twisted chiral (for N=4). The spin range in the two cases is $(0, 0) \rightarrow (N/2, 0)$ or $(0, 0) \rightarrow (N/4, N/4)$. The superfields can be multiplied in a chiral way or in a non chiral way. In the first case one reproduces a superfield of the same structure, in the latter case one gets a long multiplet with $2^{4N} r$ components, with r the (finite) dimension of some representation of $SL(2, C) \times SU(N)$. The spin range of a long multiplet is $(\Delta j_L, \Delta j_R) = (N/2, N/2)$.

An example of long multiplet is the non-chiral multiplication of two short multiplet with $j_L = j_R = 0$. This has as highest spin component a spin 4 singlet. This should be contrasted with the massless graviton multiplet obtained by tensoring, in a (twisted) chiral way, two self-conjugate singleton multiplets (with $j_L = j_R = 0$).

3. Short multiplets

Let us consider in details the case of N=4 SYM. An abelian N=4 vector multiplet corresponds to the self-conjugate representation of $SU(2, 2/4)$. The N=4 fundamental multiplet $W_{[AB]}(x, \theta, \bar{\theta})$, $A = 1, \dots, 4$ satisfies the constraints [15],

$$\begin{aligned} W_{[AB]} &= \frac{1}{2} \epsilon_{ABCD} \bar{W}_{[CD]} \\ \mathcal{D}_{\alpha A} W_{[BC]} &= \mathcal{D}_{\alpha [A} W_{BC]}. \end{aligned} \tag{3.1}$$

and contains, as first component, a set of six scalars $\phi_{[AB]}$ in the 6 of SU(4), which will be denoted also $\phi_l, l = 1, \dots, 6$. The superfield itself will be also denoted W_l .

$(x, \theta, \bar{\theta})$ can be extended to an harmonic superspace [19], where the superfield (now denoted W without indices) can be considered as a twisted chiral superfield [20]. In this way, all the product W^p are still twisted chiral superfields and therefore are short multiplets [19]. In terms of the superfield defined in eq. (3.1), we have $W^p = W_{\{l_1 \dots l_p\}}$ -traces [20].

The superfield $Tr W^2$ gives the supercurrent multiplet [14,15] and the massless graviton in AdS₅ [21]. Notice that we are taking a trace in color space in order to get gauge invariant operators. The tower of superfields $Tr W^p$ is the set of CFT composite operators in short multiplets which is in one to one correspondence with the KK states in AdS₅ $\times$ S^5 [10]. Using the explicit component expansion of W , given, for example, in [10], and performing explicitly the superfield multiplication, we obtain the full spectrum of KK states computed in [16]. The relation between masses and conformal dimensions of the CFT operators is that predicted by superconformal invariance and discussed in section 1 (formulae (1.15)-(1.20)).

We can explicitly list the operators in W_p which are in a (0,p,0) SU(4) representation and which therefore survive when fermions are neglected and only constant values of the bosonic fields $\phi_l, F_{\mu\nu}$ are retained [23].

In terms of the singleton fields $\phi_l, F_{\alpha\beta}, F_{\dot{\alpha}\dot{\beta}}$, ($F_{\alpha\beta} = \sigma_{\alpha\beta}^{\mu\nu} F_{\mu\nu}, F_{\dot{\alpha}\dot{\beta}} = \bar{F}_{\alpha\beta} = \sigma_{\dot{\alpha}\dot{\beta}}^{\mu\nu} F_{\mu\nu}$) we have,

$$Tr(\phi_{\{l_1 \dots l_p\}}) - \text{traces} \quad (0, p, 0) \quad (3.2)$$

$$Tr(\phi_{\{l_1 \dots l_{p-1}\}} F_{\alpha\beta}) - \text{traces} \quad (0, p-1, 0) \quad (3.3)$$

$$Tr(\phi_{\{l_1 \dots l_{p-2}\}} F_{\alpha\beta} F^{\alpha\beta}) - \text{traces} \quad (0, p-2, 0) \quad (3.4)$$

$$Tr(\phi_{\{l_1 \dots l_{p-2}\}} F_{\alpha\beta} F_{\dot{\alpha}\dot{\beta}}) - \text{traces} \quad (0, p-2, 0) \quad (3.5)$$

$$Tr(\phi_{\{l_1 \dots l_{p-3}\}} F_{\alpha\beta} F^{\alpha\beta} F_{\dot{\alpha}\dot{\beta}}) - \text{traces} \quad (0, p-3, 0) \quad (3.6)$$

$$\text{Tr}(\phi_{\{l_1} \cdots \phi_{l_{p-4}}\} F_{\alpha\beta} F^{\alpha\beta} F_{\dot{\alpha}\dot{\beta}} F^{\dot{\alpha}\dot{\beta}}) - \text{traces} \quad (0, p-4, 0) \quad (3.7)$$

We observe that SU(4) singlets are possible only up to $p=4$. There are exactly five of them [23]:

$$\begin{aligned} W^2 \rightarrow O_2 &= \frac{1}{2}(F^2 \pm F\tilde{F}) \quad (\tilde{F}_{\mu\nu} = \frac{i}{2}\epsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}) \\ W^2 \rightarrow T_{\mu\nu} &= F_{\mu\rho} F_{\nu\rho} - \frac{1}{4}\eta_{\mu\nu}(F_{\sigma\rho})^2 \\ W^3 \rightarrow O_3 &= F_{\mu\sigma} F_{\rho\sigma} F_{\rho\nu} - \frac{1}{4}(F_{\sigma\rho})^2 F_{\mu\nu} \\ W^4 \rightarrow O_4 &= \frac{1}{4}[(F^2)^2 - (F\tilde{F})^2] = F_{\mu\rho} F_{\nu\rho} F_{\mu\sigma} F_{\nu\sigma} - \frac{1}{4}(F^2)^2 \end{aligned} \quad (3.8)$$

In the AdS/CFT correspondence, these operators are seen, by analyzing the Born-Infeld D3 brane action, to couple to the s-wave of the type IIB (complex) dilaton, graviton, a self-dual combination of the NS-NS and R-R anti-symmetric tensors and a combination of the dilation mode of the internal (S^5) metric and the four-form anti-symmetric field with components on S^5 [24,2,25,10].

4. Long multiplets

The simplest example of long multiplet is easily constructed. By tensoring two singleton we can obtain either a spin 2 multiplet (which is again a twisted chiral superfield) or a spin 4 multiplet. The six scalars have a product which decomposes as 20+1 under SU(4). The 20 are the first components of the spin 2 multiplet which corresponds to the massless graviton in AdS₅. The singlet is the first component of a spin 4 multiplet which is not contained in the supergravity states in AdS₅, but should correspond to a massive string state [2,3].

The first component of this long multiplet is [15],

$$\text{Tr}\phi_l\phi^l : \quad \text{Tr}(W_{[AB]}W_{[CD]}\epsilon^{ABCD})|_{\theta=0} \quad (4.1)$$

and can be roughly interpreted as (the non-abelian generalization of) the radial relative positions of the D3 branes in AdS₅, while the highest one is the spin 4, made with combinations of the following operators,

$$\text{Tr}(\phi^l \mathcal{D}_{\alpha_1}^{\leftrightarrow} \cdots \mathcal{D}_{\alpha_4}^{\leftrightarrow} \phi_l), \text{Tr}(F_{\mu\rho} \mathcal{D}_{\alpha_1}^{\leftrightarrow} \mathcal{D}_{\alpha_2}^{\leftrightarrow} F_{\nu\rho}), \text{Tr}(\bar{\lambda}^A \gamma_\mu \mathcal{D}_{\alpha_1}^{\leftrightarrow} \mathcal{D}_{\alpha_2}^{\leftrightarrow} \mathcal{D}_{\alpha_3}^{\leftrightarrow} \lambda_A) - (\text{traces}) \quad (4.2)$$

with dimension $E_0 = 6$. There are also spin (1,1) with $E_0 = 4$ in the 1+15+20+... of SU(4), and spin 1 with $E_0 = 3$ in the 1+15+... of SU(4). In the free-field case, only the quoted representations appear.

This long multiplet is the N=4 embedding of the Konishi multiplet[26]. In N=1 language, the Konishi multiplet is $\Sigma = S_i e^V \bar{S}_i$, where $S_i, i = 1, \dots, 3$ are the three chiral multiplets of the N=4 theory and V is the N=1 vector superfield. Σ satisfies

$$\bar{D}\bar{D}\Sigma = W \quad (4.3)$$

where W is the superpotential multiplet $W = g\epsilon_{ijk}f_{abc}S^{ia}S^{jb}S^{kc}$ where f_{abc} are the structure constants of the gauge group. In the N=4 notations, the superpotential W is in the 10 of SU(4) since it is related to the gravitino mass term:

$$W|_{\theta=0} \rightarrow \mathcal{W}_{[AB]} = gTr\phi_{[AB]}\phi^{[BC]}\phi_{[CD]} \quad (4.4)$$

The θ^2 terms in the N=4 superfield are,

$$\theta_\alpha^A \theta_b \epsilon^{aB} \epsilon^{\alpha\beta} \mathcal{W}_{[AB]} + h.c. + \theta_\alpha^A \sigma^{\alpha\beta\mu\nu} \theta_\beta^B L_{\mu\nu} + \theta_\alpha^A \sigma_\mu^{\alpha\dot{\alpha}} \bar{\theta}_{\dot{\alpha}}^B (J_{\mu A}^B + \delta_A^B t_\mu) + \dots \quad (4.5)$$

For an unconstrained superfield these 120 terms split into $(1/2, 1/2)(15 + 1) + ((1, 0) + (0, 1))(6) + (0, 0)(10 + \bar{10})$ i.e. $120=64+36+20$. This is the θ^2 term of a scalar superfield with 2^{16} components. In the free field case, $\mathcal{W}_{[AB]} \rightarrow 0$, $\partial^\mu J_{\mu A}^B = \partial^\mu t_\mu \rightarrow 0$ and we obtain a massless multiplet with $2^8 \times 5$ components. Note that in free field theory there are additional conserved currents. This can be understood because in free N=4 Maxwell theory there is an additional SU(4) invariant fermionic current and two SU(4) currents which rotate independently scalars and fermions. In fact, in the free-field limit, this corresponds to the statement that there are infinitely many massless representations in the product of two singleton representations [12,9]. Since any irreducible representation which is contained in the product of two singletons is massless in AdS₅, in free field theory, the spin 4, 7/2, 3, 5/2, 2, 3/2, 1 would be conserved conformal fields. If the N=4 abelian multiplet is extended to a non-abelian SU(N) YM interacting multiplet, as given by eq. (4.3), then $Tr\phi_l\phi_m - \frac{1}{6}Tr\phi_p\phi^p$ is the first component of the graviton multiplet, while $Tr\phi_l\phi^l$ is the first component of a long massive spin 4 multiplet in AdS₅, which now contains all the 2^{16} components. Only in the abelian (free) case, this spin 4 multiplet become massless with only $2^8 \times 5$ physical components.

In the free-field theory limit the Konishi multiplet corresponds to the $j_L = j_R = 1$ massless spin 4 multiplet [6]. In the interacting theory, the OPE of two chiral primary operators is expected to contain also the higher spin multiplets with $j_L = j_R = s/2$ ($s=2, \dots$). These massive representations will contain massive states with maximum spin $j = j_L + j_R + 2 = s + 2$ and contain combinations of operators of the form:

$$Tr(\phi^l \mathcal{D}_{\alpha_1}^{\leftrightarrow} \cdots \mathcal{D}_{\alpha_{s+2}}^{\leftrightarrow} \phi_l), Tr(F_{\alpha_1 \rho} \mathcal{D}_{\alpha_2}^{\leftrightarrow} \cdots \mathcal{D}_{\alpha_{s+1}}^{\leftrightarrow} F_{\alpha_{s+2} \rho}), Tr(\bar{\lambda}^A \gamma_{\alpha_1} \mathcal{D}_{\alpha_2}^{\leftrightarrow} \cdots \mathcal{D}_{\alpha_{s+2}}^{\leftrightarrow} \lambda_A) - (\text{traces}) \quad (4.6)$$

These conformal operators should correspond to string states in N=4 supermultiplets up to spin $j_{MAX} = s + 2$.

In the free field theory limit, the multiplet is massless only if the dimension $l(j_L, j_R)$ of a given operator is $2 + j_L + j_R$. For other conformal dimensions the multiplet is massive. Note that, for unconstrained multiplets the conformal dimension l is an arbitrary real number satisfying $l > 2 + j_L + j_R$. This is consistent with the fact that in N=4 SYM theory a generic long multiplet has anomalous dimension, eventually related, at strong coupling, to the stringy massive excitations [2,3].

5. Conformal invariance constraints on the CFT Green functions

The correspondence between N=4 SYM and type IIB string theory on $AdS_5 \times S^5$ can be explicitly used to compute field theory Green functions in the limit in which the α' and string loop corrections can be neglected and the string theory reduces to the classical supergravity [27,28,29]. In the SYM theory this corresponds to the t'Hooft limit $N \rightarrow \infty$ and $x = g^2 N$ fixed, when x is also large [1]. The supergravity therefore describes the strong coupling dynamics of the large N limit of the N=4 SYM. All the long multiplet, which are associated to string excitations, are predicted to have large anomalous dimension $h = x^{1/4}$ [2,3], and their contribution to OPE and Green functions disappears in the strong coupling limit. In this way, at strong coupling, the OPEs and the Green functions get contribution only from the short (KK) states, whose general form has been discussed in section 3. However, when the $1/x$ (α') corrections are included, we can expect contributions also from the long multiplets.

A long multiplet contributing to the YM supercurrent Green function is just the Konishi multiplet discussed in section 4. It is indeed known that, at weak coupling, the Konishi multiplet appears in the OPE of the supercurrent multiplet [30]. Higher spin

multiplets, of the form discussed in section 4, are expected to contribute to the OPE as well. In N=1 notation, indicating with $J_{\alpha\dot{\alpha}}(z), z = (x, \theta, \bar{\theta})$ the supercurrent, we have [30]

$$\begin{aligned} J(z)J(z') &= \frac{c}{(s\bar{s})^3} + \frac{\Sigma(z')}{(s\bar{s})^{2-h/2}} + \dots \\ \Sigma(z)\Sigma(z') &= \frac{c'}{(s\bar{s})^{2+h}} + \frac{\Sigma(z')}{(s\bar{s})^{1+h/2}} + \dots \end{aligned} \quad (5.1)$$

where $s = x - x' + i\theta\gamma\bar{\theta}'$. Here $c = \frac{1}{24}(3N_v + N_\chi)$ and $c' = N_\chi$, where N_v and N_χ are the number of vectors and chiral multiplets (including color multiplicities), respectively.

This free field result for c and c' does not receive corrections up to two loops. For a generic N=1 theory with N_χ chiral multiplets in the representation T of the gauge group with a superpotential $W = Y_{ijk}\phi^i\phi^j\phi^k$, the two loop value for c is [31]

$$c = \frac{1}{24}(3N_v + N_\chi + N_v \frac{\beta(g)}{g} - \gamma_i^i) \quad (5.2)$$

where $\beta(g)$ is the one-loop beta function and γ_j^i the one-loop anomalous dimensions [32],

$$\gamma_i^j = \frac{1}{16\pi^2} \left(\frac{1}{2} Y_{ikm} Y^{jkm} - 2g^2 C_i^j(T) \right) \quad (5.3)$$

The two-loop value of c' was computed in [30] and reads

$$c' = N_\chi + 2\gamma_i^i. \quad (5.4)$$

In the N=4 SYM ($Y_{ijk} \rightarrow g f_{abc} \epsilon_{ijk}$) we see that there are no corrections up to two loop to the free-field value of c and c' . In general, c , which can be related to an R-current anomaly, can be proved to be not-renormalized at all orders [30]. However, the anomalous dimension of Σ is not zero also for conformal invariant theories and reads, at the first perturbative order,

$$h = \frac{3}{16\pi^2} \frac{Y_{ijk} Y^{ijk}}{N_\chi} \quad (5.5)$$

For N=4 we have $h = \frac{3}{16\pi^2} x$ [30], but this value is corrected, at strong coupling, to $x^{1/4}$ [2,3], and the contribution of the Konishi multiplet to the OPE, as well as of all the other higher spin long multiplets which acquire the same anomalous dimension $x^{1/4}$, becomes subleading.

³ The indices i contain both the color and the flavor indices. Similarly, T is, in general, a reducible representation of the gauge group.

We can use conformal invariance to get information on the structure of the OPE and Green functions [33]. We will give formulae valid for arbitrary space-time dimension d . Let us consider only the constraints coming from the $O(d,2)$ algebra. Supersymmetry will further imply selection rules on the operators which may appear in a given OPE expansion.

Consider, for simplicity, the OPE of two primary scalars A and B . On the hypercone, we can write the following expansion[33]:

$$A(\eta)B(\eta') = \sum_{n=0}^{\infty} (\eta \cdot \eta')^{-\frac{1}{2}(l_A+l_B-l_n+n)} D^{nA_1 \dots A_n}(\eta, \eta') O_{A_1 \dots A_n}(\eta') \quad (5.6)$$

using the pseudo-differential operator,

$$\begin{aligned} D^{nA_1 \dots A_n}(\eta, \eta', \partial') &= \eta^{A_1} \dots \eta^{A_n} D^{-\frac{1}{2}(l_A-l_B+l_n+n)}(\eta, \eta', \partial') \\ D(\eta, \eta', \partial') &= \eta \cdot \eta' \partial'^2 - 2(\eta \cdot \partial')(1 + \eta \cdot \partial') \end{aligned} \quad (5.7)$$

which is well defined when $\eta^2 = \eta'^2 = \eta \cdot \eta' = 0$.

Let us consider the case of a scalar operator O ($n=0$). Using the previous formula, the contribution of all the descendants of a given primary operator O , of dimension l , can be re-summed [33]

$$\begin{aligned} A(x)B(0) &= \left(\frac{1}{x^2}\right)^{(l_A+l_B-l)/2} \frac{\Gamma(l)}{\Gamma((l+l_A-l_B)/2)\Gamma((l-l_A+l_B)/2)} C_{AB}^O \times \\ &\int_0^1 du u^{(l_A-l_B+l)/2-1} (1-u)^{-(l_A-l_B+l)/2-1} {}_0F_1\left(l+1-d/2; -\frac{x^2}{4}u(1-u)\partial_x^2\right) O(ux) + \dots \end{aligned} \quad (5.8)$$

Here ${}_0F_1(\nu; z) = \sum_{h=0}^{\infty} \frac{1}{h!} \frac{\Gamma(\nu)}{\Gamma(\nu+h)} z^h$ is a generalized hypergeometric function.

Using the previous formula for the complete contribution of a given scalar operator O to the OPE, we can obtain the contribution of O and all its descendants to a four-point function of scalars.

Conformal invariance implies that an n -point Green's function depends on an arbitrary function of $n(n-3)/2$ parameters if

$$\frac{n(n-3)}{2} \leq nd - \frac{(d+2)(d+1)}{2} \quad (5.9)$$

and $nd-(d+2)(d+1)/2$ otherwise. The functional form of two and three-point functions is therefore completely fixed, while the four-point function depends on an arbitrary function of two conformal invariant parameters:

$$\begin{aligned} \langle 0|A(x)B(y)C(z)D(t)|0 \rangle &= [(x-y)^2]^{-l_B} [(x-z)^2]^{-\frac{1}{2}(l_A-l_B+l_C-l_D)} \times \\ &[(x-t)^2]^{-\frac{1}{2}(l_A-l_B-l_C+l_D)} [(z-t)^2]^{-\frac{1}{2}(l_C+l_D-l_A+l_B)} f(\rho, \eta) \end{aligned} \quad (5.10)$$

where

$$\rho = \frac{(x-t)^2(z-y)^2}{(x-y)^2(z-t)^2}, \quad \eta = \frac{(x-z)^2(y-t)^2}{(x-y)^2(z-t)^2}. \quad (5.11)$$

We can analyse the contribution of a conformal scalar and all its descendants to the four-point function in the s-channel, by simply using twice eq. (5.8). We obtain the formula [33]:

$$\begin{aligned} f(\rho, \eta) = & \Gamma(l) \eta^{\frac{1}{2}(l_A - l_B + l_C - l_D)} \rho^{-\frac{1}{2}(l + l_C - l_D)} \frac{\Gamma(-\frac{1}{2}(l_A - l_B + l_C - l_D))}{\Gamma((l - l_A + l_B)/2) \Gamma((l + l_A - l_B)/2)} \\ & F_4 \left(\frac{1}{2}(l + l_C - l_D), \frac{1}{2}(l + l_A - l_B); l + 1 - \frac{d}{2}; 1 + \frac{1}{2}(l_A - l_B + l_C - l_D); \frac{1}{\rho}; \frac{\eta}{\rho} \right) + \\ & \left(\frac{\rho}{\eta} \right)^{\frac{1}{2}(l_A - l_B + l_C - l_D)} [(l_A - l_B) \rightarrow -(l_A - l_B), (l_C - l_D) \rightarrow -(l_C - l_D)]. \end{aligned} \quad (5.12)$$

where F_4 is a double hypergeometric function.

For identical states and $d=4$, we obtain (up to a multiplicative constant)

$$f(\rho, \eta) = \rho^{-\frac{1}{2}} F_4 \left(\frac{l}{2}, \frac{l}{2}; l - 1, 1; \frac{1}{\rho}, \frac{\eta}{\rho} \right). \quad (5.13)$$

This result may be relevant in relating the four-point graviton amplitude in AdS_5 to the boundary correlator of four stress-energy tensors. The tree-level supergravity result should correspond, in the boundary CFT, to the exchange of chiral primary operators with canonical conformal dimension. The α' (or $1/x$) expansion [34] should receive contributions from the unknown OPE coefficients of the chiral multiplets, which cannot be specified by simply using conformal invariance, and also from the long multiplets. In the case of a four-point function, the form of the Green function is not completely specified by conformal invariance. We can determine the unknown function $f(\rho, \eta)$ in the case of a long multiplet, and confront it with the one for the exchange of a chiral multiplet, by expanding eq. (5.12) for large $l = x^{1/4}$. Note that the function $f(\rho, \eta)$ depends only on x . All the N dependence of the Green's function is encoded in the OPE coefficients.

Acknowledgements

S. F. is supported in part by the DOE under grant DE-FG03-91ER40662, Task C, the NSF grant PHY94-07194, and by the ECC Science Program SCI*-CI92-0789 (INFN-Frascati). We would like to thank D. Anselmi, L. Girardello, M. Gunaydin, D. Z. Freedman, C. Frønsdal, M. Porrati and L. Rastelli for helpful discussions. One of us (S.F.) would like to acknowledge the Institute of Theoretical Physics at Santa Barbara, where part of this work was done.

References

- [1] J. M. Maldacena, *The Large N Limit of Superconformal Field Theories and Supergravity*, hep-th/9705104.
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, *Gauge Theory Correlators from Non-Critical String Theory*, hep-th/9802109.
- [3] E. Witten, *Anti-de Sitter Space And Holography*, hep-th/9802150.
- [4] C. W. Gibbons and P. K. Townsend, Phys. Rev. Lett. 71 (1993) 3754; M. P. Blencowe and M. J. Duff, Phys. Lett. B203 (1988) 229; Nucl. Phys B310 (1988), 389; M. J. Duff, Class. Quantum Grav. 5 (1988) 189; E. Bergshoeff, M. J. Duff, C. N. Pope and E. Sezgin, Phys. Lett. B199 (1988) 69; H. Nicolai, E. Sezgin and Y. Tanii, Nucl. Phys B305 (1988) 483.
- [5] G. Mack and A. Salam, Ann. Phys. 53 (1969) 174.
- [6] M. Gunaydin, D. Minic and M. Zagermann, *4D Doubleton Conformal Theories, CPT and IIB String on $AdS_5 \times S^5$* , hep-th/9806042.
- [7] P. A. M. Dirac, Ann. Math. 37 (1936), 429.
- [8] S. Ferrara, A. F. Grillo and R. Gatto, Ann. Phys. 76 (1973) 161.
- [9] S. Ferrara and C. Frønsdal, *Conformal Maxwell theory as a singleton field theory on AdS_5 , IIB three branes and duality*, hep-th/971223; *Gauge Fields as Composite Boundary Excitations*, hep-th/9802126.
- [10] S. Ferrara, C. Frønsdal and A. Zaffaroni, *On $N=8$ Supergravity on AdS_5 and $N=4$ Superconformal Yang-Mills theory*, hep-th/9802203.
- [11] M. Flato and C. Frønsdal, J. Math. Phys. 22 (1981) 1100; Phys. Lett. B172 (1986) 412.
- [12] M. Flato and C. Frønsdal, Lett. Math. Phys 2 (1978) 421; Phys. Lett. 97B (1980) 236.
- [13] E. Angelopoulos, M. Flato, C. Frønsdal and D. Sternheimer, Phys. Rev. D23 (1981) 1278.
- [14] E. Bergshoeff, M. De Roo and B de Wit, Nucl. Phys. B182 (1981) 173.
- [15] P. Howe, K. S. Stelle and P. K. Townsend, Nucl. Phys. B192 (1981) 332.
- [16] H. J. Kim, L. J. Romans and P. van Nieuwenhuizen, Phys. Rev. D23 (1981) 1278.
- [17] D. Z. Freedman and H. Nicolai, Nucl. Phys. B237 (1984) 342.
- [18] B. Binegar, C. Frønsdal and W. Heidenreich, J. Math. Phys. 24 (1983) 2828.
- [19] P. S. Howe and P. C. West, *Non-perturbative Green's functions in theories with extended superconformal symmetry*, hep-th/9509140; Phys. Lett. B389 (1996) 273; *Is $N=4$ Yang-Mills Theory Soluble?*, hep-th/9611074; Phys.Lett. B400 (1997) 307.
- [20] L. Andrianopoli and S. Ferrara, *K-K excitations on $AdS_5 \times S^5$ as $N=4$ "primary" superfields*, hep-th/9803171.
- [21] M. Gunaydin, L. J. Romans and N. P. Warner, Phys. Lett. 154B (1985) 268; M. Pernici, K. Pilch and P. van Nieuwenhuizen, Nucl. Phys. B259 (1985) 460.

- [22] M. Gunaydin and N. Marcus, *Class. Quantum Grav.* 2 (1985) L11.
- [23] S. Ferrara, M. Lledo and A. Zaffaroni, *Born-Infeld Corrections to D3 brane Action in $AdS_5 \times S_5$ and $N=4$, $d=4$ Primary Superfields*, hep-th/9805082.
- [24] I. R. Klebanov, *Nucl.Phys.* B496 (1997) 231.
- [25] S. R. Das and S. P. Trivedi, *Three Brane Action and The Correspondence Between $N=4$ Yang Mills Theory and Anti De Sitter Space*, hep-th/9804149.
- [26] K. Konishi, *Phys. Lett.* B135 (1984) 439.
- [27] D. Z. Freedman, S. D. Mathur, A. Matusis, L. Rastelli, *Correlation functions in the $CFT(d)/AdS(d+1)$ correspondence*, hep-th/9804058.
- [28] H. Liu and A. A. Tseytlin, *$D=4$ Super Yang Mills, $D=5$ gauged supergravity and $D=4$ conformal supergravity*, hep-th/9804083.
- [29] S. Lee, S. Minwalla, M. Rangamani and N. Seiberg, *Three-Point Functions of Chiral Operators in $D=4$, $N=4$ SYM at Large N* , hep-th/9806074.
- [30] D. Anselmi, M. Grisaru, A. Johansen, *Nucl.Phys.* B491 (1997) 221; D. Anselmi, D. Z. Freedman, M. T. Grisaru, A. A. Johansen, *Phys. Lett.* B394 (1997) 329
- [31] I. Jack, *Nucl. Phys.* B253 (1985) 323.
- [32] For a recent discussion, see I. Jack, T. D. Jones and C. G. North, *Phys. Lett.* B386 (1996) 13.
- [33] S. Ferrara, A. F. Grillo and R. Gatto, *Lett. Nuovo Cimento* 2 (1971) 1363. S. Ferrara, A. F. Grillo, R. Gatto and G. Parisi, *Nucl. Phys.* B49 (1972) 77; *Nuovo Cimento* 19 (1974) 667.
- [34] T. Banks and M. B. Green, *JHEP*05(1988)002.